









Reppisch bei Dietikon

2020 Hauptmessstelle 912

Anforderung: Temp. > 10 °C = max. 0.2 mg NH₄-N/l
Temp. < 10 °C = max. 0.4 mg NH₄-N/l







Mischungstoxizität Wirkstoffgruppen

A, B	8.0	7.1	2.9	3.9	2.6	5.7	7.7	9.6	5.7	7.6	8.2	5.8	3.6	4.9	8.3	6.1	7.6	4.6	9.5	4.7	1.8	1.8	3.7	4.6	
H	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.2	0.1	0.2	0.2	0.2	0.5	0.2	0.0	0.7	0.7	0.1	0.0	0.1	0.1	0.0	0.0	0.0
F	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
INS	0.0	0.1	0.7	0.0	0.3	0.3	0.5	1.4	0.5	0.8	1.8	0.4	0.4	0.5	1.3	1.3	5.4	0.2	0.8	0.3	0.2	0.0	0.0	1.8	0.0

Mischungstoxizität taxonomische Gruppen

P	0.5	0.6	0.3	0.3	0.3	1.5	1.7	3.5	1.6	1.4	0.8	0.4	0.7	1.2	1.0	1.0	1.9	1.4	2.5	2.2	2.1	0.4	1.6	2.1	0.2	0.3
I	7.9	7.2	3.5	3.9	2.8	5.9	8.1	10	6.0	14	9.7	6.3	3.8	5.2	9.3	6.8	12	4.5	13	11	9.5	4.7	11	18	3.7	4.6
V	7.5	6.5	2.6	3.7	2.4	4.2	6.1	7.6	4.4	6.4	7.7	5.7	3.3	4.4	7.7	5.3	6.6	4.0	10	8.6	7.6	4.5	9.2	14	3.6	4.4

Jahresbeurteilung

16	Arzneimittel, Biozide, ...
1.4	Herbizide
0.1	Fungizide
14	Insektizide
3.5	Pflanzen
19.0	Invertebraten
14.4	Vertebraten